

# CAT I™

Static standing frame

Stander CAT I™ was created with a view to help children in adoption of the upright position. The upright standing contributes to the acquisition by children a new experience from the nearest environment and stimulates their psycho-physical development. In addition, it regulates the physiological processes and oxygenates the body of the child. There are many advantages during use of the CAT I™. Adjustable tilt angle (from 25° to 85°) even with a child in the device. Extensive system of belts and supports makes the child feel safe and secure. As far as this product is concerned, it is made of ecologic and environmentally friendly materials, and is also hand painted.



Easy operation

24<sup>+</sup>


24-month warranty



Stable construction



Standing frame designed with the youngest in mind.



Gas spring helps you to set the proper tilt angle of the device.



System of belts and supports makes your child feel safe and secure.



Adjustable footrests allow for smooth correction of foot positioning.



70 - 100 cm

Indicative user height



25 kg

Max. user weight



For **more details** about the product, please visit our website.

**This is a medical device.**  
**Use it in accordance with the user manual or the label.**

## INDICATIONS

Cerebral palsy (CP) – various types and forms | Brain and spinal cord injuries of various etiologies with limb paresis, severe balance disturbances, and coordination impairments | Demyelinating diseases (multiple sclerosis – MS, amyotrophic lateral sclerosis – ALS, and others) | Meningocele | Muscle diseases with paresis and paralysis (including dystrophy, myasthenia gravis, polymyositis) | Genetic syndromes with limb paresis | Degenerative and metabolic diseases of the nervous system | Degenerative and rheumatic joint diseases | Other diseases with limb paresis, paralysis, and musculoskeletal impairments | After injuries, following traumatic brain injuries or spinal cord injuries with limb paresis and paralysis, particularly with quadriplegia (as part of the rehabilitation process or as permanent orthopedic support)

## CONTRAINDICATIONS

**Observation:** Monitor user during standing. Stop therapy if paleness, sweating, fainting, joint overload, or body deformations occur; consult with a doctor or physiotherapist | **Absolute Contraindication:** Critical condition of the user. | **Relative Contraindications:** Resting hypertension > 200/110 mmHg, pulmonary hypertension, heart rhythm disorders, unstable asthma, unstable metabolic issues, moderate heart valve disease. | Conditions like healing fractures or severe osteoporosis that prevent weight-bearing | Incomplete fracture healing (unless approved by a doctor) | Postoperative joint load restrictions | Pain from dislocation, excessive tissue stretch, or pressure; dislocation without pain is not contraindicated | Significant hip/knee flexion or ankle contractures; ensure no excessive pressure or stretch on affected areas.