DALMATIAN INVENTOTM

Multifunctional stander

DALMATIAN Invento™ is a versatile rehabilitation chair suitable for both children and young people. It serves as a chair for adopting a sitting position and transitions into a static standing frame for rehabilitation, with a lying function throughout the process. Equipped with a gas spring, the chair allows therapists and caregivers to easily position the child upright without removing them from the chair. The standing angle can be adjusted from 0° to 90°, and it features a safety vest and pelvic belt with the Hold & Pull™ system for quick and secure stabilization. Made from eco-friendly materials and hand-painted, DALMATIAN Invento™ is designed to support the rehabilitation process effectively.



Tilt-in-space function



24-month warranty



Stable construction



Easy operation

Colors

ATMO™ upholsteryBreathable and comfortable




Grass



Sea



Lying position



Stander position



100 - 165 cm Indicative user height



25 - 60 kg Max. user weight



For **more details** about the product, please visit our website.

This is a medical device.

Use it in accordance with the user manual or the label.

INDICATIONS

Cerebral palsy (CP) – various types and forms | Brain and spinal cord injuries of various etiologies with limb paresis, severe balance disturbances, and coordination impairments | Demyelinating diseases (multiple sclerosis – MS, amyotrophic lateral sclerosis – ALS, and others) | Meningocele | Muscle diseases with paresis and paralysis (including dystrophy, myasthenia gravis, polymyositis) | Genetic syndromes with limb paresis | Degenerative and metabolic diseases of the nervous system | Degenerative and rheumatic joint diseases | Other diseases with limb paresis, paralysis, and musculoskeletal impairments | After injuries, following traumatic brain injuries or spinal cord injuries with limb paresis and paralysis, particularly with quadriplegia (as part of the rehabilitation process or as permanent orthopedic support)

CONTRAINDICATIONS

Observation: Monitor user during standing. Stop therapy if paleness, sweating, fainting, joint overload, or body deformations occur; consult with a doctor or physiotherapist | Absolute Contraindication: Critical condition of the user. | Relative Contraindications: Resting hypertension > 200/110 mmHg, pulmonary hypertension, heart rhythm disorders, unstable asthma, unstable metabolic issues, moderate heart valve disease. | Conditions like healing fractures or severe osteoporosis that prevent weight-bearing | Incomplete fracture healing (unless approved by a doctor) | Postoperative joint load restrictions | Pain from dislocation, excessive tissue stretch, or pressure; dislocation without pain is not contraindicated | Significant hip/knee flexion or ankle contractures; ensure no excessive pressure or stretch on affected areas.