

KIDOO™ is a therapeutic chair that is perfect for therapy and fun as well as other daily activities such as learning or taking meals. The device can be used at home, school, kindergarten or in a group of peers. KIDOO™ turns out to be an indispensable tool in the rehabilitation process, both in terms of therapy and social integration. This orthopedic chair provides the child with proper support and safety, moreover, it can be easily adjusted to meet your child's needs and dysfunctions. Pelvic belt height adjustable seat and armrests further enhance comfort and safety of the user.



Lightweight



24-month warranty



Stable construction



Tilt-in-space function

Colors

MEDICUS™ upholstery

Waterproof and stream-permable



Med Sea



Med Skin



Med Red



Med Lilac



Med Green



Med Gray

EXTREME-MED™ upholstery

Stain-resistant and wear-resistant



Pillarbox



Medical upholstery used in this product is vapor permeable, waterproof and easy to disinfect.



Seat depth adjustment and backrest tilt angle available in standard equipment.




90 - 160 cm

Indicative user height



20 - 70 kg

Max. user weight



For **more details** about the product, please visit our website.

This is a medical device.
Use it in accordance with the user manual or the label.

INDICATIONS

Cerebral palsy (CP) – various types and forms | Brain and spinal cord injuries of various etiologies with limb paresis, severe balance disturbances, and coordination impairments | Demyelinating diseases (multiple sclerosis – MS, amyotrophic lateral sclerosis – ALS, and others) | Meningocele | Muscle diseases with paresis and paralysis (including dystrophy, myasthenia gravis, polymyositis) | Genetic syndromes with limb paresis | Degenerative and metabolic diseases of the nervous system | Degenerative and rheumatic joint diseases | Other diseases with limb paresis, paralysis, and musculoskeletal impairments | After injuries, following traumatic brain injuries or spinal cord injuries with limb paresis and paralysis, particularly with quadriplegia (as part of the rehabilitation process or as permanent orthopedic support)

CONTRAINDICATIONS

Observation: Monitor user during standing. Stop therapy if paleness, sweating, fainting, joint overload, or body deformations occur; consult with a doctor or physiotherapist | **Absolute Contraindication:** Critical condition of the user. | **Relative Contraindications:** Resting hypertension > 200/110 mmHg, pulmonary hypertension, heart rhythm disorders, unstable asthma, unstable metabolic issues, moderate heart valve disease. | Conditions like healing fractures or severe osteoporosis that prevent weight-bearing | Incomplete fracture healing (unless approved by a doctor) | Postoperative joint load restrictions | Pain from dislocation, excessive tissue stretch, or pressure; dislocation without pain is not contraindicated | Significant hip/knee flexion or ankle contractures; ensure no excessive pressure or stretch on affected areas.